

$U_q(N)$ GAUGE THEORIES

Leonardo Castellani

*Istituto Nazionale di Fisica Nucleare
and
Dipartimento di Fisica Teorica
Via P. Giuria 1, I-10125 Torino, Italy***Abstract**

Improving on an earlier proposal, we construct the gauge theories of the quantum groups $U_q(N)$. We find that these theories are consistent also with an ordinary (commuting) spacetime. The bicovariance conditions of the quantum differential calculus are essential in our construction. The gauge potentials and the field strengths are q -commuting “fields”, and satisfy q -commutation relations with the gauge parameters. The transformation rules of the potentials are given explicitly, and generalize the ordinary infinitesimal gauge variations. The q -lagrangian invariant under the $U_q(N)$ variations has the Yang-Mills form $F_{\mu\nu}^i F_{\mu\nu}^j g_{ij}$, the “quantum metric” g_{ij} being a generalization of the Killing metric.

DFTT-74/92
December 1992

e-mail addresses: 31890::castellani, castellani@torino.infn.it

In recent times there have been some attempts [1] to construct gauge theories of quantum groups [2]. These theories are continuously connected with ordinary Yang-Mills theories, just as quantum groups are continuously connected with ordinary Lie groups. In ref. [3] we have proposed a geometric approach to the construction of q -gauge theories, based on the differential calculus on q -groups developed in [4-9], and we have found a q -lagrangian invariant under the action of $[SU(2) \times U(1)]_q$. Here we generalize the procedure to any quantum group $U_q(N)$.

Spacetime is taken to be ordinary (commutative) spacetime, but the whole discussion holds for a generic q -spacetime.

We start by recalling some essential facts about quantum Lie algebras. There are many ways to deform Lie algebras: here we consider the one that naturally comes from the quantum differential calculus, i.e. the q -algebra of the quantum tangent vectors. The notations will be as in ref. [9].

The q -algebra we obtain from a bicovariant differential calculus is given by the q -commutations between the quantum generators T_i :

$$T_i T_j - \Lambda^{kl}_{ij} T_k T_l = \mathbf{C}_{ij}{}^k T_k \quad (1)$$

The braiding matrix Λ and the q -structure constants $\mathbf{C}_{ij}{}^k$ can be expressed in terms of the R -matrix of the corresponding q -group, as shown in [6] and further discussed in [7-9], and satisfy four conditions:

$$\Lambda^{ij}_{kl} \Lambda^{lm}_{sp} \Lambda^{ks}_{qu} = \Lambda^{jm}_{kl} \Lambda^{ik}_{qs} \Lambda^{sl}_{up} \quad (\text{Yang - Baxter equation}) \quad (2a)$$

$$\mathbf{C}_{mi}{}^r \mathbf{C}_{rj}{}^n - \Lambda^{kl}_{ij} \mathbf{C}_{mk}{}^r \mathbf{C}_{rl}{}^n = \mathbf{C}_{ij}{}^k \mathbf{C}_{mk}{}^n \quad (q - \text{Jacobi identities}) \quad (2b)$$

$$\mathbf{C}_{is}{}^j \Lambda^{sq}_{rl} \Lambda^{ir}_{pk} + \mathbf{C}_{rl}{}^q \Lambda^{jr}_{pk} = \Lambda^{jq}_{ri} \Lambda^{si}_{kl} \mathbf{C}_{ps}{}^r + \Lambda^{jq}_{pi} \mathbf{C}_{kl}{}^i \quad (2c)$$

$$\Lambda^{ir}_{mk} \Lambda^{ks}_{nl} \mathbf{C}_{rs}{}^j = \Lambda^{ij}_{kl} \mathbf{C}_{mn}{}^k \quad (2d)$$

cf. for ex. [4,5,9]. The last two conditions are trivial in the limit $q \rightarrow 1$ ($\Lambda^{ij}_{kl} = \delta_l^i \delta_k^j$). The q -Jacobi identity shows that the matrix $(T_i)_j{}^k \equiv \mathbf{C}_{ji}{}^k$ is a representation (the adjoint representation) of the q -algebra (1).

This is all the information we need from the general theory of [4]. We introduce now the definition of field strength

$$F_{\mu\nu} \equiv \frac{1}{2}(\partial_\mu A_\nu - \partial_\nu A_\mu + A_\mu A_\nu - A_\nu A_\mu) = \partial_{[\mu} A_{\nu]} - A_{[\mu} A_{\nu]} \quad (3)$$

where

$$A_\mu \equiv A_\mu^i T_i$$

The gauge potentials A_μ^i satisfy the q -commutations:

$$A_{[\mu}^i A_{\nu]}^j = -\frac{1}{q^2 + q^{-2}}(\Lambda + \Lambda^{-1})^{ij}{}_{kl} A_{[\mu}^k A_{\nu]}^l \quad (4)$$

and simply commute with the quantum generators T_i . The square parentheses around the μ, ν indices stand for ordinary antisymmetrization. The inverse Λ^{-1} of the braiding matrix always exists [4] and is defined by $(\Lambda^{-1})^{ij}{}_{kl} \Lambda^{kl}{}_{mn} = \delta_m^i \delta_n^j$. The rule (4) is inspired by the q -commutations of exterior products of left-invariant one-forms on the quantum groups $U_q(N)$:

$$A^i \wedge A^j = -\frac{1}{q^2 + q^{-2}}(\Lambda + \Lambda^{-1})^{ij}{}_{kl} A^k \wedge A^l \quad (5)$$

deduced in ref. [9]. Then replacing

$$A^i = A_\mu^i dx^\mu \quad (6)$$

into eq. (5) yields the rule (4). Note that the differentials dx^μ are ordinary spacetime differentials, commuting with A_μ^i .

For future use we define the projectors:

$$P_S^{ij}{}_{kl} \equiv \frac{1}{2 + q^2 + q^{-2}}[(q^2 + q^{-2}) + (\Lambda + \Lambda^{-1})]^{ij}{}_{kl} \quad (7a)$$

$$P_A^{ij}{}_{kl} \equiv \frac{1}{2 + q^2 + q^{-2}}[2I - (\Lambda + \Lambda^{-1})]^{ij}{}_{kl} \quad (7b)$$

where $I^{ij}{}_{kl} \equiv \delta_k^i \delta_l^j$ is the identity matrix. It is not difficult to verify the properties:

$$P_S P_S = P_A P_A = I, \quad P_S P_A = P_A P_S = 0 \quad (8)$$

due to the spectral decomposition of Λ [7]:

$$(\Lambda + q^2 I)(\Lambda + q^{-2} I)(\Lambda - I) = 0, \quad (9)$$

a consequence of the Hecke relation for the $U_q(N)$ R -matrix [2c].

We are now ready to compute the explicit form of the field strength:

$$\begin{aligned} F_{\mu\nu}^i T_i &= \partial_{[\mu} A_{\nu]}^i T_i + A_{[\mu}^i A_{\nu]}^j T_i T_j = \\ &= \partial_{[\mu} A_{\nu]}^i T_i + \alpha A_{[\mu}^i A_{\nu]}^j T_i T_j + (1 - \alpha) A_{[\mu}^i A_{\nu]}^j T_i T_j = \\ &= \partial_{[\mu} A_{\nu]}^i T_i + \alpha A_{[\mu}^i A_{\nu]}^j T_i T_j - \frac{(1 - \alpha)}{q^2 + q^{-2}} (\Lambda + \Lambda^{-1})^{kl}{}_{ij} A_{[\mu}^i A_{\nu]}^j T_k T_l \\ &= \partial_{[\mu} A_{\nu]}^i T_i + \alpha A_{[\mu}^i A_{\nu]}^j [T_i T_j - \frac{(1 - \alpha)}{\alpha(q^2 + q^{-2})} (\Lambda + \Lambda^{-1})^{kl}{}_{ij} T_k T_l] \end{aligned} \quad (10)$$

where we have used the q -commutations (4). Setting $\alpha = \frac{2}{2 + q^2 + q^{-2}}$ we find:

$$\begin{aligned}
F_{\mu\nu}^i T_i &= \partial_{[\mu} A_{\nu]}^i T_i + \frac{2}{2+q^2+q^{-2}} A_{[\mu}^i A_{\nu]}^j [T_i T_j - \frac{1}{2}(\Lambda + \Lambda^{-1})^{kl}{}_{ij} T_k T_l] \\
&= \partial_{[\mu} A_{\nu]}^i T_i + \frac{1}{2+q^2+q^{-2}} A_{[\mu}^i A_{\nu]}^j [\mathbf{C}_{ij}{}^n - (\Lambda^{-1})^{kl}{}_{ij} \mathbf{C}_{kl}{}^n] T_n
\end{aligned} \tag{11}$$

since

$$T_i T_j - \frac{1}{2}(\Lambda + \Lambda^{-1})^{kl}{}_{ij} T_k T_l = \frac{1}{2}[\mathbf{C}_{ij}{}^n - (\Lambda^{-1})^{kl}{}_{ij} \mathbf{C}_{kl}{}^n] T_n \tag{12}$$

(use (1) and the relation one obtains after multiplying (1) by Λ^{-1}). Let us define

$$\mathbf{C}_{kl}{}^n \equiv C_{kl}{}^n - \Lambda^{ij}{}_{kl} C_{ij}{}^n \tag{13}$$

Then

$$\mathbf{C}_{ij}{}^n - (\Lambda^{-1})^{kl}{}_{ij} \mathbf{C}_{kl}{}^n = (2+q^2+q^{-2}) P_A{}^{kl}{}_{ij} C_{kl}{}^n \tag{14}$$

and the field strength can finally be written as:

$$F_{\mu\nu}^i = \partial_{[\mu} A_{\nu]}^i + P_A{}^{kl}{}_{mn} C_{kl}{}^i A_{[\mu}^m A_{\nu]}^n \tag{15}$$

Note that the Cartan-Maurer equations for the left invariant one-forms on the quantum groups $U_q(N)$ can be written as:

$$dA^i + P_A{}^{kl}{}_{mn} C_{kl}{}^i A^m \wedge A^n = 0$$

so that the right-hand side of eq. (13) is to be expected (see ref.[9] for a discussion on quantum Cartan-Maurer equations and q -curvatures).

We next define the gauge variations:

$$\delta A_\mu = -\partial_\mu \varepsilon - A_\mu \varepsilon + \varepsilon A_\mu \tag{16}$$

with

$$\varepsilon \equiv \varepsilon^i T_i$$

and postulate the commutations

$$\varepsilon^i A_\mu^j = \Lambda^{ij}{}_{kl} A_\mu^k \varepsilon^l \tag{17}$$

Under the variations (16) the field strength transforms as:

$$\delta F_{\mu\nu} = \varepsilon F_{\mu\nu} - F_{\mu\nu} \varepsilon \tag{18}$$

Indeed the calculation is identical to the usual one.

Using (17) the gauge variations of A_μ^i become

$$\delta A_\mu^i = -\partial_\mu \varepsilon^i - A_\mu^j \varepsilon^k \mathbf{C}_{jk}{}^i \tag{19}$$

Let us derive the commutations of ε^i with $F_{\mu\nu}^j$. To be as general as possible we rewrite the q -commutations (17) as:

$$\varepsilon^i A^j = \Lambda^{ij}_{kl} A^k \varepsilon^l \quad (20)$$

i.e. we do not use (6): our results will be valid even for q -spacetime. We prove now that:

$$\varepsilon^i F^j = \Lambda^{ij}_{kl} F^k \varepsilon^l \quad (21)$$

Taking the exterior derivative of (20) yields (we omit wedge symbols in the following):

$$(d\varepsilon^i)A^j + \varepsilon^i dA^j = \Lambda^{ij}_{kl} [(dA^k)\varepsilon^l + A^k(d\varepsilon^l)] \quad (22)$$

Requesting that the $d\varepsilon$ terms separately cancel leads to:

$$(d\varepsilon^i)A^j = \Lambda^{ij}_{kl} A^k (d\varepsilon^l) \quad (23a)$$

$$\varepsilon^i dA^j = \Lambda^{ij}_{kl} (dA^k) \varepsilon^l \quad (23b)$$

Eq. (23b) tells us that the dA^i part of F^i does satisfy (21). What happens to the quadratic piece ? We should find that

$$\varepsilon^i A^m A^n [\mathbf{C}_{mn}^j - (\Lambda^{-1})^{pq}_{mn} \mathbf{C}_{pq}^j] = \Lambda^{ij}_{kl} A^m A^n [\mathbf{C}_{mn}^k - (\Lambda^{-1})^{pq}_{mn} \mathbf{C}_{pq}^k] \varepsilon^l \quad (24)$$

in order to satisfy (21). The left-hand side can be ordered as $AA\varepsilon$ by using (20) twice, and comparing it to the right-hand side we find an equation of the type $\Lambda\Lambda\mathbf{C} - \Lambda\Lambda\Lambda^{-1}\mathbf{C} = \Lambda\mathbf{C} - \Lambda\Lambda^{-1}\mathbf{C}$. The first and the third term cancel because of the bicovariance condition (2d); the second and the fourth term give:

$$\Lambda^{ir}_{mk} \Lambda^{ks}_{nl} (\Lambda^{-1})^{pq}_{rs} = \Lambda^{ij}_{kl} (\Lambda^{-1})^{pq}_{mn} \mathbf{C}_{pq}^k \quad (25)$$

Multiplying both sides by Λ^{mn}_{uv} and using the Yang-Baxter equation (2b) leads again to the bicovariance condition (2d). Thus the ε, F commutations (21) hold because of the bicovariance conditions (2b) and (2d), and can be used in eq. (18) to yield:

$$\delta F_{\mu\nu}^i = -F_{\mu\nu}^j \varepsilon^k \mathbf{C}_{jk}^i$$

Does the condition (2c) also play a role ? The answer is yes: it ensures that the commutation relations (4) (or equivalently (5)) are preserved under the q -gauge transformations (16). Indeed the gauge variation of (5) is:

$$\delta A^i A^j + A^i \delta A^j = -\frac{1}{q^2 + q^{-2}} (\Lambda + \Lambda^{-1})^{ij}_{kl} (\delta A^k A^l + A^k \delta A^l) \quad (26)$$

Substitute the variations given by (16) and consider first the terms without derivatives, after having them reordered as $AA\varepsilon$ via eq. (20). Using (5) on the left-hand side leads exactly to the bicovariance condition (2c). The terms in (26) with the $d\varepsilon$ derivatives can

be reordered as $A(d\varepsilon)$ using (23a), and are easily seen to cancel because of the cubic relation (9).

The commutations between F^i and A^j can be obtained by taking the exterior derivative of the A, A commutations (5):

$$F^i A^j - A^i F^j = -\frac{1}{q^2 + q^{-2}} (\Lambda + \Lambda^{-1})^{ij}_{kl} (F^k A^l - A^k F^l) \quad (27)$$

Indeed the terms trilinear in A that arise after using $dA = R - CAA$ do cancel, since they cancel in the case $F^i = 0$, i.e. for left-invariant A^i , and the wedge products are unaltered in the case $F^i \neq 0$ (For a discussion on the “softening” of quantum group geometry, see ref. [9]).

Finally, we construct the q -lagrangian invariant under the $U_q(N)$ quantum Lie algebra. We set:

$$\mathcal{L} = F_{\mu\nu}^i F_{\mu\nu}^j g_{ij} \quad (28)$$

where g_{ij} , the q -analogue of the Killing metric, is determined by requiring the invariance of $\mathcal{L}$ under the q -gauge transformations (16). Under these transformations, the variation of the lagrangian (28) is given by

$$\delta\mathcal{L} = -\mathbf{C}_{mn}^i F_{\mu\nu}^m \varepsilon^n F_{\mu\nu}^j g_{ij} - \mathbf{C}_{mn}^j F_{\mu\nu}^i F_{\mu\nu}^m \varepsilon^n g_{ij} \quad (29)$$

After using (21) for reordering the terms as $FF\varepsilon$ we find that $\delta\mathcal{L}$ vanishes when

$$\mathbf{C}_{mn}^i \Lambda^{nj}_{rs} g_{ij} + \mathbf{C}_{rs}^j g_{mj} = 0 \quad (30)$$

We have not yet succeeded in finding the general solution for the q -Killing metric g_{ij} . However, eq. (30) is not difficult to solve in particular cases. For example, in the $U_q(2) \equiv [SU(2) \times U(1)]_q$ case of ref. [3], we find that

$$g_{ij} = \begin{pmatrix} -2(1 - q^2) + 2r & -2 & 0 & 0 \\ -2 & 2r & 0 & 0 \\ 0 & 0 & 0 & 2q^2(r + 1) \\ 0 & 0 & 2q^2(r + 1) & 0 \end{pmatrix} \quad (31)$$

is the most general q -metric satisfying (30), depending on the deformation parameter q and on an extra arbitrary parameter r , related to the ratio of the $SU(2)$ and $U(1)$ coupling constants (see ref. [3]). For $q \rightarrow 1$, $r \rightarrow 1$ one recovers the usual Killing metric of $SU(2) \times U(1)$.

Note 1: It is clear that g_{ij} should generalize the $q = 1$ expression for the Killing metric $g_{ij} = \text{Tr}(\mathbf{C}_i \mathbf{C}_j) = \mathbf{C}_{ri}^s \mathbf{C}_{sj}^r$. A quantum trace in the adjoint representation is needed.

Note 2: The results of this Letter could be extended to the quantum groups of the B, C, D , series.

References

- [1] A.P. Isaev and Z. Popowicz, *q-Trace for the Quantum Groups and q-deformed Yang-Mills Theory*, Wroclaw preprint ITP UWr 786-91; I.Y. Aref'eva and I.V. Volovich, Mod. Phys. Lett. **A6** (1991) 893; Phys. Lett. **B264** (1991) 62; S. Meljanac and S. Pallua, *Classical field lagrangian and deformed algebras*, DFPF/92/TH/15; A. Dimakis and F. Müller-Hoissen, *Noncommutative differential calculus, gauge theory and gravitation*, GOET-TP-33/92; K. Wu and R.-J. Zhang, Commun. Theor. Phys. **17** (1992) 175; T. Brzeziński and S. Majid, *Quantum group gauge theory on quantum spaces*, DAMTP/92-27 and *Quantum group gauge theory on classical spaces*, DAMTP/92-51.
- [2a] V. Drinfeld, Sov. Math. Dokl. **32** (1985) 254.
- [2b] M. Jimbo, Lett. Math. Phys. **10** (1985) 63; **11** (1986) 247.
- [2c] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Algebra and Analysis, **1** (1987) 178.
- [2d] S. Majid, Int. J. Mod. Phys. **A5** (1990) 1.
- [3] L. Castellani, Phys. Lett. **B292** (1992) 93.
- [4] S.L. Woronowicz, Publ. RIMS, Kyoto Univ., Vol. **23**, 1 (1987) 117; Commun. Math. Phys. **111** (1987) 613 and Commun. Math. Phys. **122**, (1989) 125.
- [5] D. Bernard, *Quantum Lie algebras and differential calculus on quantum groups*, Proc. 1990 Yukawa Int. Seminar, Kyoto; Phys. Lett. **260B** (1991) 389.
- [6] B. Jurčo, Lett. Math. Phys. **22** (1991) 177.
- [7] U. Carow-Watamura, M. Schlieker, S. Watamura and W. Weich, Commun. Math. Phys. **142** (1991) 605.
- [8] B. Zumino, *Introduction to the Differential Geometry of Quantum Groups*, LBL-31432 and UCB-PTH-62/91, notes of a plenary talk given at the 10-th IAMP Conf., Leipzig (1991); F. Müller-Hoissen, J. Phys. A **25** (1992) 1703; X.D. Sun and S.K. Wang, *Bicovariant differential calculus on quantum group $GL_q(n)$* , Worldlab-Beijing preprint CCAST-92-04, ASIAM-92-07, ASITP-92-12 (1992); A. Sudbery, Phys. Lett. **B284** (1992) 61; P. Schupp, P. Watts and B. Zumino, *Differential Geometry on Linear Quantum Groups*, preprint LBL-32314, UCB-PTH-92/13 (1992) and *Bicovariant quantum algebras and quantum Lie algebras*, LBL-32315, UCB-PTH-92/14; B. Zumino, *Differential calculus on quantum spaces and quantum groups*, LBL-33249, UCB-PTH-92/41.
- [9] P. Aschieri and L. Castellani, *An introduction to non-commutative differential geometry on quantum groups*, preprint CERN-TH.6565/92, DFTT-22/92 (1992), to be publ. in Int. Jou. Mod. Phys. A.